

# FIFRA Section 24(c) Special Local Need



FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA



(EPA Reg. No. 59639-99)  
EPA SLN No. NC-150007

FOR CONTROL OF WINTER ANNUAL BROADLEAF WEEDS IN CLARY SAGE

**This label is valid until December 31, 2025 or until otherwise amended, withdrawn, cancelled or suspended.**

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered container product label for Valor® SX Herbicide.

This labeling must be in the possession of the user at the time of application.

**FOR WEED CONTROL IN CLARY SAGE GROWN FOR THE ESSENCE  
AND ESSENTIAL OIL INDUSTRY  
(Do not use treated clary sage for food or feed)**

## **PREPLANT APPLICATION FOR RESIDUAL CONTROL OF WINTER ANNUAL BROADLEAF WEEDS IN CLARY SAGE**

- Apply 1.0 to 2.0 oz/A of *Valor* SX Herbicide as a preplant application when applied 30 days before planting clary sage with soils that contain less than 80% sand.
- **If soils contain greater than 80% sand**, apply 1 oz/A *Valor* SX Herbicide 30 days before planting clary sage or 2.0 oz/A 45 days before planting.
- When tank mixing, use 1.0 oz/A for enhanced burndown and limited residual weed control.
- Use the 2.0 oz/A rate for residual weed control through planting.
- Apply 3.0 oz/A of *Valor* SX Herbicide as a preplant application when applied 45 or more days before planting clary sage.
- Sequential preplant applications are prohibited.

## **Application Restrictions**

- Treated soil that is splashed onto newly emerged sage may result in temporary crop injury.
- A minimum of 0.5 inch of rainfall/irrigation must pass between application and planting of clary sage.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- Apply in a minimum of 15 GPA of water to ensure uniform soil coverage and adequate coverage of emerged weeds.
- Do not apply more than 3.0 oz/A of *Valor* SX Herbicide preplant.
- Do not apply more than 7.0 oz of *Valor* SX Herbicide per acre per season.

**POSTEMERGENCE APPLICATION FOR RESIDUAL CONTROL OF WINTER ANNUAL  
BROADLEAF WEEDS IN CLARY SAGE**

- Apply 1.0 to 2.0 oz/A of *Valor* SX Herbicide as a postemergence application when clary sage is a minimum of 4 inches in diameter.

**Application Restrictions**

- *Valor* SX Herbicide can be applied postemergence to clary sage up to the time of flower bud initiation.
- Do not apply *Valor* SX Herbicide to clary sage after it begins flower bud initiation.
- Do not apply more than 2 oz of *Valor* SX Herbicide per acre during a single application.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- Do not tank mix *Valor* SX Herbicide with any adjuvant, fertilizer or pest control product or severe injury to clary sage will occur.
- Apply in a minimum of 15 GPA of water to ensure adequate coverage of emerged weeds.
- Do not apply more than 7.0 oz of *Valor* SX Herbicide per acre per season.

**Please contact Valent U.S.A. Corporation at 1-800-6-VALENT (682-5368) if you have questions regarding use of this product.**

**FIFRA 24(c) Registrant:** Valent U.S.A. LLC  
P.O. Box 5075  
San Ramon, CA 94583

*Valor* is a registered trademark of Valent U.S.A. LLC.