

SePRO Corporation 11550 N. Meridian St., Suite 600, Carmel, IN 46032 USA

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NORTH CAROLINA

Only for use by North Carolina Department of Transportation and their authorized users.



Preemergence herbicide for use in sulphur cosmos (*Cosmos sulphureus*) and zinnia (*Zinnia* sp.) plantings to control glyphosate-resistant Palmer amaranth.

Active Ingredient

Fluridone:	
1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone	
Other Ingredients	
Total	
Equivalent to 1.2 pounds fluridone per gallon.	

EPA Reg. No. 67690-78

EPA SLN No. NC-190001

This label for Brake expires and must not be distributed or used in accordance with this SLN registration after December 31, 2028.

ATTENTION

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.
- Refer to the Brake container label, EPA Reg. No. 67690-78, for complete Directions for Use and all applicable restrictions and precautions.
- This supplemental label and the entire Brake container label must be in the possession of the user at the time of pesticide application.

DIRECTIONS FOR USE

Preemergence Applications

Apply this product as a preemergence treatment to coarse, medium or fine textured soils at rates from 11 to 21 fluid ounces per acre. **If applying less than 16 fluid ounces per acre, another residual herbicide effective at controlling the target weed species must be tank mixed with this product.** Use in accordance with the most restrictive label limitations and precautions of the products used in the tank-mix. Do not exceed any labeled rate and only tank mix with other herbicides labeled for use in sulphur cosmos and zinnia. For best results, apply behind the planter (i.e. at planting) or within 36 hours after planting. Do not apply more than 21 fluid ounces per acre in a single application. Moisture is necessary to activate this product in soil for residual weed control. Dry weather following applications of this product may reduce effectiveness. For improved weed control, a minimum of 0.5 inches of rain or overhead irrigation is recommended following application.

Use Restrictions

- Only for use by North Carolina Department of Transportation and their authorized users.
- **DO NOT** apply more than 32 fluid ounces of this product per acre per year (equivalent to 0.3 lb fluridone per acre).
- Apply prior to planting or within 36 hours after planting.
- DO NOT apply this product over the top of plants as injury will likely occur.

Minor chlorosis and stunting of sulphur cosmos and zinnia may occur, especially if heavy rainfall occurs during or soon after emergence, but plants typically outgrow these effects and develop normally. User should assume these risks before using this product.

Brake is a registered trademark of SePRO Corporation. FPL20190322 © Copyright 2019 SePRO Corporation