

FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Staple LX[®]

EPA Reg. No. 352-613

EPA SLN No. NC 200001

For Distribution and Use Only By The State of North Carolina
Department of Transportation (NCDOT)

CONTROL OF CERTAIN BROADLEAF AND GRASS WEEDS IN WILDFLOWERS

Expiration date: This label expires and must not be distributed or used in accordance with this SLN registration after **July 31, 2026**.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Special Conditions and Risks of Use

USE OF Staple LX herbicide (THE "PRODUCT") ON WILDFLOWERS (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;

(5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of North Carolina.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Preemergence Application – Apply Staple® LX preemergence to wildflowers at 1.3 - 2.1 oz/acre.

Wildflower species shown to have acceptable tolerance to Staple LX preemergence application include Bachelors Button, Black-Eye Susan, Ox-eye Daisy, Plains Coreopsis, Shasta Daisy, Sulphur Cosmos, and Zinnia.

Wildflower species with unacceptable tolerance to Staple LX preemergence application include Baby Blue Eyes, Bird's Eye, California Poppy, Catchfly, Clasping Coneflower, Corn Poppy, Dame's Rocket, Gloriosa Daisy, Indian Blanket, Lance-leaved Coreopsis, Mountain Garland, Purple Coneflower, Rocket Larkspur, Toadflax, and Wallflower.

Postemergence Application – Apply Staple LX postemergence to wildflowers at 2.6 oz/acre. A nonionic surfactant (0.25% v/v) is recommended to improve control.

Wildflower species shown to have acceptable tolerance to Staple LX postemergence application include Bachelors Button, Gloriosa Daisy, Ox-eye Daisy, Plains Coreopsis, Rocket Larkspur, Shasta Daisy, Sweet William, and Zinnia.

Wildflower species with unacceptable tolerance to Staple LX postemergence application include Baby Blue Eyes, Bird's Eye, Black-eyed Susan, California Poppy, Catchfly, Clasping Coneflower, Corn Poppy, Dame's Rocket, Gloriosa Daisy, Indian Blanket, Lance-leaved Coreopsis, Mountain Garland, Purple Coneflower, Rocket Larkspur, Sulphur Cosmos, Toadflax, and Wallflower.

Staple LX may be applied as a preemergence and postemergence application to other tolerant wildflower species that are supported by research at North Carolina State University.

™®Trademarks of Corteva Agriscience and its affiliated companies

R640-002

Approved: ___/___/___

Initial printing/Replaces R645-001